



BlueCross BlueShield of Oklahoma

If a conflict arises between a Clinical Payment and Coding Policy (“CPCP”) and any plan document under which a member is entitled to Covered Services, the plan document will govern. If a conflict arises between a CPCP and any provider contract pursuant to which a provider participates in and/or provides Covered Services to eligible member(s) and/or plans, the provider contract will govern. “Plan documents” include, but are not limited to, Certificates of Health Care Benefits, benefit booklets, Summary Plan Descriptions, and other coverage documents. BCBSOK may use reasonable discretion interpreting and applying this policy to services being delivered in a particular case. BCBSOK has full and final discretionary authority for their interpretation and application to the extent provided under any applicable plan documents.

Providers are responsible for submission of accurate documentation of services performed. Providers are expected to submit claims for services rendered using valid code combinations from Health Insurance Portability and Accountability Act (“HIPAA”) approved code sets. Claims should be coded appropriately according to industry standard coding guidelines including, but not limited to: Uniform Billing (“UB”) Editor, American Medical Association (“AMA”), Current Procedural Terminology (“CPT®”), CPT® Assistant, Healthcare Common Procedure Coding System (“HCPCS”), ICD-10 CM and PCS, National Drug Codes (“NDC”), Diagnosis Related Group (“DRG”) guidelines, Centers for Medicare and Medicaid Services (“CMS”) National Correct Coding Initiative (“NCCI”) Policy Manual, CCI table edits and other CMS guidelines.

Claims are subject to the code edit protocols for services/procedures billed. Claim submissions are subject to claim review including but not limited to, any terms of benefit coverage, provider contract language, medical policies, clinical payment and coding policies as well as coding software logic. Upon request, the provider is urged to submit any additional documentation.

Prostate Biopsies

Policy Number: CPCPLAB005

Version 1.0

Enterprise Clinical Payment and Coding Policy Committee Approval Date: July 5, 2023

Plan Effective Date: November 1, 2023

Description

BCBSOK has implemented certain lab management reimbursement criteria. Not all requirements apply to each product. Providers are urged to review Plan documents for eligible coverage for services rendered.

Reimbursement Information:

1. In the initial diagnosis of prostate cancer as a follow-up to abnormal PSA results, presence of a palpable nodule on digital rectal examination, or suspicious radiologic findings, prostate biopsy, or

comprehensive 3D (three-dimensional) mapping with biopsy involving 12 core extended sampling (See NOTE 1 below) **may be reimbursable**.

2. When the clinical suspicion of prostate cancer remains in an individual for whom an initial biopsy was negative for prostate cancer, follow-up biopsy (excluding prostate saturation biopsy) **may be reimbursable**.
3. Prostate saturation biopsy or comprehensive 3D mapping with biopsy **are not reimbursable** for the diagnosis, staging and management of prostate cancer.

Note 1: One vial per sextant, with no more than two core samples per vial.

Procedure Codes

The following is not an all-encompassing code list. The inclusion of a code does not guarantee it is a covered service or eligible for reimbursement.

Codes
0443T, 55700, 55706, 88305, G0416

References:

Balducci, L., Pow-Sang, J., Friedland, J., & Diaz, J. I. (1997). Prostate cancer. *Clin Geriatr Med*, 13(2), 283-306. <http://dx.doi.org/>

Bekelman, J. E., Rumble, R. B., Chen, R. C., Pisansky, T. M., Finelli, A., Feifer, A., Nguyen, P. L., Loblaw, D. A., Tagawa, S. T., Gillissen, S., Morgan, T. M., Liu, G., Vapiwala, N., Haluschak, J. J., Stephenson, A., Touijer, K., Kungel, T., & Freedland, S. J. (2018). Clinically Localized Prostate Cancer: ASCO Clinical Practice Guideline Endorsement of an American Urological Association/American Society for Radiation Oncology/Society of Urologic Oncology Guideline. *Journal of Clinical Oncology*, 36(32), 3251-3258. <https://doi.org/10.1200/JCO.18.00606>

Bell, K. J., Del Mar, C., Wright, G., Dickinson, J., & Glasziou, P. (2015). Prevalence of incidental prostate cancer: A systematic review of autopsy studies. *Int J Cancer*, 137(7), 1749-1757. <https://doi.org/10.1002/ijc.29538>

Benway, B., & Andriole, G. (2021). Prostate biopsy. <https://www.uptodate.com/contents/prostate-biopsy>

Bjurlin, M. A., Carroll, P. R., Eggener, S., Fulgham, P. F., Margolis, D. J., Pinto, P. A., Rosenkrantz, A. B., Rubenstein, J. N., Rukstalis, D. B., Taneja, S. S., & Turkbey, B. (2020). Update of the Standard Operating Procedure on the Use of Multiparametric Magnetic Resonance Imaging for the Diagnosis, Staging and Management of Prostate Cancer. *J Urol*, 203(4), 706-712. <https://doi.org/10.1097/ju.0000000000000617>

Brown, L. C., Ahmed, H. U., Faria, R., El-Shater Bosaily, A., Gabe, R., Kaplan, R. S., Parmar, M., Collaco-Moraes, Y., Ward, K., Hindley, R. G., Freeman, A., Kirkham, A., Oldroyd, R., Parker, C., Bott, S., Burns-Cox, N., Dudderidge, T., Ghei, M., Henderson, A., . . . Emberton, M. (2018). Multiparametric MRI to improve detection of prostate cancer compared with transrectal ultrasound-guided prostate biopsy alone: the PROMIS study. *Health Technol Assess*, 22(39), 1-176. <https://doi.org/10.3310/hta22390>

Coakley, F. V., Oto, A., Alexander, L. F., Allen, B. C., Davis, B. J., Froemming, A. T., Fulgham, P. F., Hosseinzadeh, K., Porter, C., Sahni, V. A., Schuster, D. M., Showalter, T. N., Venkatesan, A. M., Verma, S., Wang, C. L., Remer, E. M., & Eberhardt, S. C. (2017). ACR Appropriateness Criteria((R)) Prostate Cancer- Pretreatment Detection, Surveillance, and Staging. *J Am Coll Radiol*, 14(5s), S245-s257. <https://doi.org/10.1016/j.jacr.2017.02.026>

Eichler, K., Hempel, S., Wilby, J., Myers, L., Bachmann, L. M., & Kleijnen, J. (2006). Diagnostic value of systematic biopsy methods in the investigation of prostate cancer: a systematic review. *J Urol*, 175(5), 1605-1612. [https://doi.org/10.1016/s0022-5347\(05\)00957-2](https://doi.org/10.1016/s0022-5347(05)00957-2)

Gosselaar, C., Roobol, M. J., Roemeling, S., Wolters, T., van Leenders, G. J., & Schroder, F. H. (2008). The value of an additional hypoechoic lesion-directed biopsy core for detecting prostate cancer. *BJU Int*, 101(6), 685-690. <https://doi.org/10.1111/j.1464-410X.2007.07309.x>

Hodge, K. K., McNeal, J. E., Terris, M. K., & Stamey, T. A. (1989). Random systematic versus directed ultrasound guided transrectal core biopsies of the prostate. *J Urol*, 142(1), 71-74; discussion 74-75. <http://dx.doi.org/>

Kantoff, P., Taplin, M.-E., & Smith, J. (2020). *Clinical presentation and diagnosis of prostate cancer - UpToDate*. <https://www.uptodate.com/contents/clinical-presentation-and-diagnosis-of-prostate-cancer>

Klotz, L., Chin, J., Black, P. C., Finelli, A., Anidjar, M., Bladou, F., Mercado, A., Levental, M., Ghai, S., Chang, S. D., Milot, L., Patel, C., Kassam, Z., Moore, C., Kasivisvanathan, V., Loblaw, A., Kebabdjian, M., Earle, C. C., Pond, G. R., & Haider, M. A. (2021). Comparison of Multiparametric Magnetic Resonance Imaging-Targeted Biopsy With Systematic Transrectal Ultrasonography Biopsy for Biopsy-Naive Men at Risk for Prostate Cancer: A Phase 3 Randomized Clinical Trial. *JAMA Oncol*, 7(4), 534-542. <https://doi.org/10.1001/jamaoncol.2020.7589>

NCCN. (2022a). NCCN Clinical Practice Guidelines in Oncology; Prostate Cancer. v2. https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf

NCCN. (2022b). NCCN Clinical Practice Guidelines in Oncology; Prostate Cancer Early Detection. v2. Retrieved June 29, 2021, from https://www.nccn.org/professionals/physician_gls/pdf/prostate_detection.pdf

Parker, C., Castro, E., Fizazi, K., Heidenreich, A., Ost, P., Procopio, G., Tombal, B., & Gillessen, S. (2020). Prostate Cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology*. <https://doi.org/10.1016/j.annonc.2020.06.011>

Pepe, P., & Aragona, F. (2007). Saturation prostate needle biopsy and prostate cancer detection at initial and repeat evaluation. *Urology*, 70(6), 1131-1135. <https://doi.org/10.1016/j.urology.2007.07.068>

Pepe, P., Garufi, A., Priolo, G. D., Galia, A., Fraggetta, F., & Pennisi, M. (2018). Is it Time to Perform Only Magnetic Resonance Imaging Targeted Cores? Our Experience with 1,032 Men Who Underwent Prostate Biopsy. *J Urol*, 200(4), 774-778. <https://doi.org/10.1016/j.juro.2018.04.061>

Pepe, P., Pennisi, M., & Fraggetta, F. (2020). How Many Cores Should be Obtained During Saturation Biopsy in the Era of Multiparametric Magnetic Resonance? Experience in 875 Patients Submitted to Repeat Prostate Biopsy. *Urology*, 137, 133-137. <https://doi.org/10.1016/j.urology.2019.11.016>

Samir S. Taneja, M. C., Marc A. Bjurlin, DO, H. Ballentine Carter, MD, Michael S. Cookson, MD, MMHC, Leonard G. Gomella, MD, FACS, David F. Penson, MD, MPH, Paul Schellhammer, MD, Steven Schlossberg MD, MBA, Dean Troyer, MD. (2015). *American Urological Association - Optimal Techniques of Prostate Biopsy and Specimen Handling* <http://www.auanet.org/guidelines/prostate-biopsy-and-specimen-handling>

Sidana, A., Watson, M. J., George, A. K., Rastinehad, A. R., Vourganti, S., Rais-Bahrami, S., Muthigi, A., Maruf, M., Gordetsky, J. B., Nix, J. W., Merino, M. J., Turkbey, B., Choyke, P. L., Wood, B. J., & Pinto, P. A. (2018). Fusion prostate biopsy outperforms 12-core systematic prostate biopsy in patients with prior negative systematic biopsy: A multi-institutional analysis. *Urol Oncol*, 36(7), 341.e341-341.e347. <https://doi.org/10.1016/j.urolonc.2018.04.002>

Siegel, R. L., Miller, K. D., & Jemal, A. (2020). Cancer statistics, 2020. *CA: A Cancer Journal for Clinicians*, 70(1). <https://doi.org/10.3322/caac.21551>

Smith, R. A., Andrews, K. S., Brooks, D., Fedewa, S. A., Manassaram-Baptiste, D., Saslow, D., Brawley, O. W., & Wender, R. C. (2018). Cancer screening in the United States, 2018: A review of current American Cancer Society guidelines and current issues in cancer screening. *CA: A Cancer Journal for Clinicians*, 68(4), 297-316. <https://doi.org/10.3322/caac.21446>

Smith, R. A., Andrews, K. S., Brooks, D., Fedewa, S. A., Manassaram-Baptiste, D., Saslow, D., & Wender, R. C. (2019). Cancer screening in the United States, 2019: A review of current American Cancer Society guidelines and current issues in cancer screening. *CA: A Cancer Journal for Clinicians*, 69(3), 184-210. <https://doi.org/10.3322/caac.21557>

Tabayoyong, W., & Abouassaly, R. (2015). Prostate Cancer Screening and the Associated Controversy. *Surg Clin North Am*, 95(5), 1023-1039. <https://doi.org/10.1016/j.suc.2015.05.001>

Thompson, J. E., Hayen, A., Landau, A., Haynes, A. M., Kalapara, A., Ischia, J., Matthews, J., Frydenberg, M., & Stricker, P. D. (2015). Medium-term oncological outcomes for extended vs saturation biopsy and transrectal vs transperineal biopsy in active surveillance for prostate cancer. *BJU Int*, 115(6), 884-891. <https://doi.org/10.1111/bju.12858>

USPSTF. (2018). Screening for prostate cancer: US Preventive Services Task Force recommendation statement. *JAMA*, 319(18), 1901-1913. <https://doi.org/10.1001/jama.2018.3710>

Wolf, A. M., Wender, R. C., Etzioni, R. B., Thompson, I. M., D'Amico, A. V., Volk, R. J., Brooks, D. D., Dash, C., Guessous, I., Andrews, K., DeSantis, C., & Smith, R. A. (2010). American Cancer Society guideline for the early detection of prostate cancer: update 2010. *CA Cancer J Clin*, 60(2), 70-98. <https://doi.org/10.3322/caac.20066>

Zaytoun, O. M., Moussa, A. S., Gao, T., Fareed, K., & Jones, J. S. (2011). Office based transrectal saturation biopsy improves prostate cancer detection compared to extended biopsy in the repeat biopsy population. *J Urol*, 186(3), 850-854. <https://doi.org/10.1016/j.juro.2011.04.069>

Policy Update History:

7/5/2023	Document updated with literature review. Reimbursement statement 1 edited for clarity. Added #2: When the clinical suspicion of prostate cancer remains in an individual for whom an initial biopsy was negative for prostate cancer, follow-up biopsy (excluding prostate saturation biopsy) may be reimbursable. References revised.
11/1/2022	New policy